

Recombinant Human IP-10/CXCL10

Information

Gene ID	
Accession #	
Alternate Names	
Source	Escherichia coli.
M.Wt	Approximately 8.6 kDa, a single non-glycosylated polypeptide chain containing 77 amino acids.
AA Sequence	VPLSRTVRCT CISISNQPVN PRSLEKLEII PASQFCPRVE IATMKKKGE KRCLNPESKA IKNLLKAVSK EMSKRSP
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 50 mM NaCl..
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human peripheral blood T-lymphocytes is in a concentration range of 10- 50 ng/ml.
Shipping Condition	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	5µg	100µg	500µg
Recombinant Human Ciliary Neurotrophic Factor	5µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

Purity	> 97 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 1 EU/μg of rHuIP-10/CXCL10 as determined by LAL method

Description

CXCL10 also known as IP-10 is belonging to the CXC chemokine family. It is encoded by the CXCL10 gene. CXCL10 was originally identified in monocytes, endothelial cells and fibroblasts as a responder to IFN- γ . This chemokine elicits its effects by binding to the cell surface chemokine receptor CXCR3. CXCL10 has been shown to be a chemoattractant for activated T lymphocytes and monocytes/macrophages. It also has other roles, such as promotion of T cell adhesion to endothelial cells, and inhibition of bone marrow colony formation and angiogenesis. The mouse homologue of human IP-10, Crg-2, has been cloned and shown to share approximately 67 % amino acid sequence identity with human IP-10.

Reference

APExBIO Technology
www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

