

Recombinant Human Fibroblast Growth Factor 17

Information

Gene ID	8822
Accession #	O60258
Alternate Names	
Source	Escherichia coli.
M.Wt	Approximately 22.6 kDa, a single non-glycosylated polypeptide chain containing 195 amino acids.
AA Sequence	MTQGENHPSP NFNQYVRDQG AMTDQLSRRQ IREYQLYSRT SGKHVQVTGR RISATAEDGN KFAKLIVETD TFGSRVRIKG AESEKYICMN KRGKLGKPS GSKKDCVFTE IVLENNYTAF QNARHEGWFM AFTRQGRPRQ ASRSRQNRQRE AHFIKRLYQG QLPPFNHAEK QKQFEFVGSA PTRRTKRTRR PQPLT
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The ED as determined by a cell proliferation assay using murine balb/c 3T3 cells is less than 10 ng/ml, corresponding to a specific activity of >1.0 × 10 IU/mg.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	5µg	100µg	500µg
Recombinant Human Fibroblast Growth Factor 17	5µg	100µg	500µg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles

- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

Purity	> 95 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 0.1 EU/μg of rHuFGF-17 as determined by LAL method.

Description

Reference



APExBIO Technology
www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

