

Recombinant Murine Lymphotoctin/XCL1

Information

Gene ID	16963
Accession #	P47993
Alternate Names	
Source	Escherichia coli.
M.Wt	Approximately 10.1 kDa, a single non-glycosylated polypeptide chain containing 93 amino acids.
AA Sequence	VGTEVLEESS CVNLQTQRLP VQKIKTYIIW EGAMRAVIFV TKRGLKICAD PEAKWVKAAL KTVDGRASTR KNMAETVPTG AQRSTSTAIT LTG
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
Formulation	Lyophilized from a 0.2 μm filtered concentrated solution in PBS, pH 7.4.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Fully biologically active when compared to standard. The ED as determined by a chemotaxis bioassay using human XCR1 transfected murine BaF3 cells is less than 100 ng/ml, corresponding to a specific activity of > 1.0 × 10 IU/mg.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	5μg	100μg	500μg
Recombinant Murine Lymphotoctin/XCL1	5μg	100μg	500μg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles


- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

Purity	> 97 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 0.1 EU/ μ g of rMuLymphotactin/XCL1 as determined by LAL method.

Description

Reference



APEX BIO Technology
www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com