

Recombinant Human TP53-induced Glycolysis and Apoptosis Regulator

Information

Gene ID	57103
Accession #	Q9NQ88
Alternate Names	
Source	Escherichia coli.
M.Wt	Approximately 30.1 kDa, a single non-glycosylated polypeptide chain containing 270 amino acids.
AA Sequence	MARFALTVVR HGETRFNKEK IIQQQGVDEP LSETGFKQAA AAGIFLNNVK FTHAFSSDLM RTKQTMHGIL ERSKFCKDMT VKYDSRLRER KYGVVEGKAL SELRAMAKAA REECPVFTPP GGETLDQVKM RGIDFFFLC QLILKEADQK EQFSQGPSN CLETSLAEIF PLGKNHSSKV NSDSGIPGLA ASVLVSHGA YMRSLFDYFL TDLKCSLPAT LSRSELSVT PNTGMSLFII NFEEGREVKP TVQCICMNLQ DHLNGLTETR
Appearance	Sterile Filtered White lyophilized (freeze-dried) powder.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles - 12 months from date of receipt, -20 to -70 °C as supplied - 1 month, 2 to 8 °C under sterile conditions after reconstitution - 3 months, -20 to -70 °C under sterile conditions after reconstitution
Formulation	Lyophilized from a 0.2 μm filtered concentrated solution in 20 mM Tris-HCl, pH8.5, 150 mM NaCl.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Biological Activity	Data not available.
Shipping Condition	Gel pack.
Handling	Centrifuge the vial prior to opening.
Usage	For Research Use Only! Not to be used in humans.

Components and Storage

Components	5μg	100μg	500μg
Recombinant Human TP53-induced Glycolysis and Apoptosis Regulator	5μg	100μg	500μg

Use a manual defrost freezer and avoid repeated freeze-thaw cycles


- 12 months from date of receipt, -20 to -70 °C as supplied
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 3 months, -20 to -70 °C under sterile conditions after reconstitution

Quality Control

Purity	> 95 % by SDS-PAGE and HPLC analyses.
Endotoxin	Less than 0.1 EU/ μ g of rHuTIGAR as determined by LAL method.

Description

Reference



APExBIO Technology
www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com